

Comparison of two methods for the definition of the infinite time limit of the time evolution operator in scattering theory

Gábor Zsolt Tóth

*International School for Advanced Studies (SISSA),
Via Beirut 2-4, 34014 Trieste, Italy*¹

*INFN, Sezione di Trieste, Italy*¹

*Research Institute for Particle and Nuclear Physics,
Hungarian Academy of Sciences, Pf. 49, 1525 Budapest, Hungary*

e-mail: tgzs@cs.elte.hu

Abstract

A comparison of the methods of averaging and adiabatic switching for the definition of the infinite time limit of the time evolution operator in scattering theory is presented. The focus is on certain basic features which are relevant mainly in quantum field theory. Specifically, the differences between the unitarity properties of the operators obtained using the two methods, the difference between the necessary energy renormalizations and the existence of Lippmann-Schwinger equations are discussed. The manifestation of these differences in perturbation expansions is also described. An illustration on the case of the ϕ^4 model using perturbation theory is presented.

Keywords: S-matrix; Hamiltonian perturbation theory; adiabatic switching; Abelian limit; self-energy corrections; renormalization

¹Until 31 October 2009.

1 Introduction

The $T \rightarrow \pm\infty$ limits of the time evolution operator $U(0, -T)$ have an important role in scattering theory and also in some other areas of quantum physics. However, in general it is not completely obvious how to define these limits. In this paper we discuss certain basic aspects of this problem; we compare two methods for the definition of $\lim_{T \rightarrow \pm\infty} U(0, -T)$, which are mentioned in the literature as the method of averaging and the method of adiabatic switching. We emphasize certain features of the two methods and highlight certain significant differences between them. The discussion is done in the context of scattering theory, where the time evolution operator is used to produce the *in* and *out* states.

In section 2 we introduce briefly the *in* and *out* states, the S-matrix elements and the time evolution operator. In sections 3 and 4 we describe the method of averaging and the method of adiabatic switching, respectively.

The main differences between the two methods are the following. 1) In general a renormalization of the free particle Hamiltonian operator is necessary, but the amounts of energy renormalizations that are needed are different for the two methods. 2) The $\lim_{T \rightarrow \pm\infty} U(0, -T)$ operators obtained by averaging are in general not unitary, therefore normalization factors have to be included in the formulas which give the *in* and *out* states. Such normalization factors are not needed in the case of the adiabatic switching. 3) The Lippmann-Schwinger equations which hold in the case of the method of averaging do not hold in the case of the method of adiabatic switching.

The difference between the two methods can be studied in perturbation theory as well. In particular, it can be seen that the two methods are identical to first order, but generally there are differences at higher orders.

The differences that we discuss are relevant quite generally, in particular also for quantum field theory; nevertheless in quantum mechanical potential scattering the differences between the two methods usually vanish.

In addition to the sections 2, 3 and 4 described above, the paper contains section 5, where we present an illustration of the ideas described in sections 3 and 4 in the case of the ϕ^4 model, and section 6, where the conclusions are presented. Appendix A contains formulas relevant for section 5.

We note that we were partially motivated to write this paper by recent development in scattering theory [1]-[13], and that this paper is related to an earlier paper [14] of ours.

2 The *in* and *out* states

Let H be the total Hamiltonian operator that describes a certain scattering process. We assume that H takes the form $H = H_0 + gH_{\text{int}}$, where H_0 is a Hamiltonian operator that describes free particles, gH_{int} is an interaction term and g is a coupling constant that can

be used as a variable for power series expansion.

The S-matrix elements are scalar products of *in* and *out* states:

$$S_{wv} = \langle w, out | v, in \rangle. \quad (1)$$

An important property of the *in* and *out* states is that they are eigenstates of H .

In quantum mechanics standard formulas (see [15, 16, 17]) that give the *in* and *out* states in terms of H , H_0 and the eigenstates of H_0 are the following:

$$|v, in\rangle = \lim_{T \rightarrow \infty} U(0, -T)|v\rangle, \quad (2)$$

$$|v, out\rangle = \lim_{T \rightarrow \infty} U(0, T)|v\rangle. \quad (3)$$

In these formulas $U(t_2, t_1)$ is the time evolution operator

$$U(t_2, t_1) = e^{\frac{i}{\hbar} H_0 t_2} e^{-\frac{i}{\hbar} H(t_2 - t_1)} e^{-\frac{i}{\hbar} H_0 t_1}, \quad (4)$$

and $|v\rangle$ is an eigenstate of H_0 . The *in* and *out* states $|v, in\rangle$ and $|v, out\rangle$ thus correspond to the eigenstate $|v\rangle$ of H_0 , and $\lim_{g \rightarrow 0} |v, in\rangle = \lim_{g \rightarrow 0} |v, out\rangle = |v\rangle$. The energy of an eigenstate $|v\rangle$ will be denoted by E_v^0 : $H_0|v\rangle = E_v^0|v\rangle$. The energy of $|v, in\rangle$ and $|v, out\rangle$ will be denoted by E_v : $H|v, in\rangle = E_v|v, in\rangle$, $H|v, out\rangle = E_v|v, out\rangle$. The formulas (2) and (3) imply that $E_v = E_v^0$ [15].

In the following sections we aim to write (2) and (3) in a form that is valid generally, in particular also in quantum field theory. We shall see that certain modifications of (2) and (3) are necessary for this, and these modifications depend on the way the $T \rightarrow \infty$ limit is defined. We will discuss two methods, which we mentioned in the introduction, for defining this limit. The equation (1) will not be changed.

We will write three different modified versions of (2) and (3); the first one is (20) and (21) (or, with some regularizations explicitly denoted, (29) and (30)), the second one is (48) and (49) (or (53) and (54)), and the third one is (55) and (56). The first version is based on the method of averaging, whereas the second and third versions are based on the method of adiabatic switching. We mention the third version only briefly, since it was the subject of [14].

3 The method of averaging

In order to properly define the $T \rightarrow \infty$ limit of $U(0, -T)|v\rangle$ and $U(0, T)|v\rangle$ an averaging can be introduced in the following way (see e.g. [18]):

$$\hat{U}_\epsilon(0, -T) = \epsilon \int_0^T d\tau e^{-\epsilon\tau} U(0, -\tau), \quad (5)$$

$$\hat{U}_\epsilon(0, T) = \epsilon \int_0^T d\tau e^{-\epsilon\tau} U(0, \tau), \quad (6)$$

and then

$$|v, in\rangle = \lim_{\epsilon \rightarrow +0} \lim_{T \rightarrow \infty} \hat{U}_\epsilon(0, -T)|v\rangle, \quad (7)$$

$$|v, out\rangle = \lim_{\epsilon \rightarrow +0} \lim_{T \rightarrow \infty} \hat{U}_\epsilon(0, T)|v\rangle. \quad (8)$$

The $T \rightarrow \infty$ limit defined using this average is also known as the Abelian limit.

The above definition for the *in* and *out* states is usually satisfactory in potential scattering (at least for sufficiently short range and regular potentials). However, in general (in quantum field theory, for example) one encounters divergences in the $\epsilon \rightarrow +0$ limit if one uses the definitions (7) and (8) and applies perturbation theory to calculate the components of the *in* and *out* states and the S-matrix elements.

These divergences occur in any theory in which the energy of (the actual) $|v, in\rangle$ and $|v, out\rangle$ states, which are eigenstates of H , is shifted with respect to the energy of $|v\rangle$ by any nonzero amount. These shifts are called self-energy corrections. We use the notation δE_v for them (thus $\delta E_v = E_v - E_v^0$). We stress that the mentioned divergences are distinct from the ultraviolet divergences which often occur in quantum field theory, and also from the type of infrared divergences which are caused by soft or collinear massless particles in theories like quantum electrodynamics and quantum chromodynamics. We shall not discuss these types of divergences in this paper. In quantum field theory the divergences that we discuss are associated with disconnected vacuum bubble diagrams and with radiative corrections on external lines. A detailed explanation can be found e.g. in section 4.6 of [19].

The problem of self-energy corrections can also be illustrated in a nonperturbative framework; for this purpose let us consider the case when the Hilbert space is finite dimensional and the spectrum of H_0 is nondegenerate. Although the scattering is trivial in such a system, the self-energy corrections are generally nonzero. We consider the generic case when H is such that for sufficiently small but nonzero values of g all eigenvalues of H are shifted from the values that they take at $g = 0$. We assume that $|v, in\rangle$, for an eigenvector $|v\rangle$ of H_0 , is defined by (7). At $g = 0$ we have $|v, in\rangle = |v\rangle$, of course. For small values of g , the limits in (7) exist, but the shift of the eigenvalues of H has the consequence that $\langle W|v, in\rangle = 0$ for any eigenvector $|W\rangle$ of H , and thus $|v, in\rangle = 0$. This can be seen in the following way. Let E_W be the eigenvalue for $|W\rangle$. Then we have

$$\langle W|U(0, -T)|v\rangle = \langle W|v\rangle \exp\left[\frac{i}{\hbar}(E_v^0 - E_W)T\right], \quad (9)$$

which is an oscillating function of T since $E_v^0 \neq E_W$. It is easy to verify that the averaging procedure yields zero for this function. This holds for all eigenvectors of H , and these eigenvectors form a basis in the Hilbert space, therefore the limits in (7) exist and $|v, in\rangle = 0$. This implies, in particular, that $|v, in\rangle$ is discontinuous at $g = 0$ as a function of g .

3.1 Renormalization of H_0

The standard solution to the problem described above is the renormalization of H_0 , which consists in replacing the initial H_0 operator in the definition of $U(t_2, t_1)$ by a new one, while H remains unchanged. $U(t_2, t_1)$ is thus replaced in (5) and (6) by

$$U^{\text{ren}}(t_2, t_1) = e^{\frac{i}{\hbar} H_0^{\text{ren}} t_2} e^{-\frac{i}{\hbar} H(t_2 - t_1)} e^{-\frac{i}{\hbar} H_0^{\text{ren}} t_1} \quad (10)$$

and we have the formulas

$$\hat{U}_\epsilon^{\text{ren}}(0, -T) = \epsilon \int_0^T d\tau e^{-\epsilon\tau} U^{\text{ren}}(0, -\tau), \quad (11)$$

$$\hat{U}_\epsilon^{\text{ren}}(0, T) = \epsilon \int_0^T d\tau e^{-\epsilon\tau} U^{\text{ren}}(0, \tau), \quad (12)$$

$$|v, in\rangle = \lim_{\epsilon \rightarrow +0} \lim_{T \rightarrow \infty} \hat{U}_\epsilon^{\text{ren}}(0, -T) |v^{\text{ren}}\rangle, \quad (13)$$

$$|v, out\rangle = \lim_{\epsilon \rightarrow +0} \lim_{T \rightarrow \infty} \hat{U}_\epsilon^{\text{ren}}(0, T) |v^{\text{ren}}\rangle. \quad (14)$$

The vector $|v^{\text{ren}}\rangle$ appearing in (13) and (14) should be an eigenvector of H_0^{ren} . We denote the eigenvalue for $|v^{\text{ren}}\rangle$ by E_v^{ren} : $H_0^{\text{ren}} |v^{\text{ren}}\rangle = E_v^{\text{ren}} |v^{\text{ren}}\rangle$. E_v^{ren} and $\Delta E_v = E_v^{\text{ren}} - E_v^0$ are called renormalized energy and energy renormalization, respectively. H_0^{ren} has to have the same eigenvalue on $|v^{\text{ren}}\rangle$ as H has on $|v, in\rangle$ and $|v, out\rangle$, i.e. $E_v^{\text{ren}} = E_v$ has to hold.

(More generally, one can replace the pair (H_0, H) by a suitable renormalized pair $(H_0^{\text{ren}}, H^{\text{ren}})$, and in this case it is also possible that $H_0 = H_0^{\text{ren}}$. For the subject of this paper it has no significance whether H is changed or not.)

In perturbation theory the condition that the $\epsilon \rightarrow +0$ limit in (13) and (14) should be convergent imposes restrictions on the H_0^{ren} operator and on the $|v^{\text{ren}}\rangle$ vectors at every order.

In [18, 20, 21] the following convenient choice is proposed:

$$H_0^{\text{ren}} = H_0 + \int di \Delta E_i |v_i\rangle \langle v_i| = \int di E_i^{\text{ren}} |v_i\rangle \langle v_i| \quad (15)$$

and

$$|v_i^{\text{ren}}\rangle = |v_i\rangle, \quad (16)$$

where the $|v_i\rangle$ vectors are eigenvectors of H_0 and constitute an orthonormal basis i.e.

$$\langle v_i | v_j \rangle = \delta(i, j), \quad (17)$$

where $\delta(i, j)$ is a Dirac-delta. The integration in (15) is understood in a general sense; it includes, for instance, summation over discrete states. The Dirac-delta is also a generalized

one. The index i labels all eigenvectors of H_0 . The ΔE_i are the energy renormalizations and $E_i^{\text{ren}} = E_i^0 + \Delta E_i$ are the renormalized energies. For all E_i^{ren} the equation

$$E_i^{\text{ren}} = E_i \quad (18)$$

has to hold, where the E_i is the eigenvalue of the H operator on $|v_i, \text{in}\rangle$ and $|v_i, \text{out}\rangle$. The renormalized energies E_i^{ren} can be calculated order by order in perturbation theory from the condition that the $\epsilon \rightarrow +0$ limit in (13) and (14) should be convergent.

The choice above for H_0^{ren} and $|v_i^{\text{ren}}\rangle$ is not suitable in some cases (we refer the reader to section 3.4 for more details), nevertheless we conjecture that it is suitable for a large class of quantum field theories.

3.2 Normalization of the *in* and *out* states

The formulas (13) and (14) are still unsatisfactory because $\lim_{\epsilon \rightarrow +0} \lim_{T \rightarrow \infty} \hat{U}_\epsilon^{\text{ren}}(0, -T)$ and $\lim_{\epsilon \rightarrow +0} \lim_{T \rightarrow \infty} \hat{U}_\epsilon^{\text{ren}}(0, T)$ are generally not unitary (more precisely, not scalar product preserving), even though $U^{\text{ren}}(t_2, t_1)$ is unitary. This is caused by the property of the averaging procedure that it assigns 0 in the $T \rightarrow \infty$ limit to functions which oscillate around 0.

In order to illustrate the nonunitarity of $\lim_{\epsilon \rightarrow +0} \lim_{T \rightarrow \pm\infty} \hat{U}_\epsilon^{\text{ren}}(0, -T)$ we consider again the case when the Hilbert space is finite dimensional and the spectrum of H_0 is nondegenerate. We assume that H_0^{ren} takes the form (15) and g is sufficiently small so that the spectrum of H is also nondegenerate. Considering the definition of $U^{\text{ren}}(0, -T)$ and the property that $|v, \text{in}\rangle$ is an eigenvector of H , we obviously have $\langle v, \text{in} | U^{\text{ren}}(0, -T) | v \rangle = \langle v, \text{in} | v \rangle$. This implies that

$$\langle v, \text{in} | v, \text{in} \rangle = \langle v, \text{in} | \lim_{\epsilon \rightarrow +0} \lim_{T \rightarrow \infty} \hat{U}_\epsilon^{\text{ren}}(0, -T) | v \rangle = \langle v, \text{in} | v \rangle. \quad (19)$$

However, $\langle v, \text{in} | v, \text{in} \rangle = \langle v, \text{in} | v \rangle$ cannot hold if $\langle v, \text{in} | v, \text{in} \rangle = \langle v | v \rangle$ unless $|v, \text{in}\rangle = |v\rangle$, which is not the case generally. Therefore $\langle v, \text{in} | v, \text{in} \rangle \neq \langle v | v \rangle$. In fact, the operator $\lim_{\epsilon \rightarrow +0} \lim_{T \rightarrow \infty} \hat{U}_\epsilon^{\text{ren}}(0, -T)$ projects $|v\rangle$ onto the (one dimensional) space spanned by $|v, \text{in}\rangle$, implying in particular that $\langle v, \text{in} | v, \text{in} \rangle < \langle v | v \rangle$.

On the basis of the above considerations, we can say that the correct form of (7) and (8) should be

$$|v, \text{in}\rangle = \lim_{\epsilon \rightarrow +0} \lim_{T \rightarrow \infty} \frac{1}{\sqrt{Z_{v, \epsilon, \text{in}}(-T)}} \hat{U}_\epsilon^{\text{ren}}(0, -T) | v^{\text{ren}} \rangle, \quad (20)$$

$$|v, \text{out}\rangle = \lim_{\epsilon \rightarrow +0} \lim_{T \rightarrow \infty} \frac{1}{\sqrt{Z_{v, \epsilon, \text{out}}(T)}} \hat{U}_\epsilon^{\text{ren}}(0, T) | v^{\text{ren}} \rangle, \quad (21)$$

where

$$Z_{v,\epsilon,in}(-T) = \frac{\langle v^{\text{ren}} | \hat{U}_\epsilon^{\text{ren}}(0, -T)^\dagger \hat{U}_\epsilon^{\text{ren}}(0, -T) | v^{\text{ren}} \rangle}{\langle v^{\text{ren}} | v^{\text{ren}} \rangle}, \quad (22)$$

$$Z_{v,\epsilon,out}(T) = \frac{\langle v^{\text{ren}} | \hat{U}_\epsilon^{\text{ren}}(0, T)^\dagger \hat{U}_\epsilon^{\text{ren}}(0, T) | v^{\text{ren}} \rangle}{\langle v^{\text{ren}} | v^{\text{ren}} \rangle}. \quad (23)$$

The numbers $Z_{v,\epsilon,in}$ and $Z_{v,\epsilon,out}$ are real and are included in (20) and (21) in order to maintain the correct normalization of $|v, in\rangle$ and $|v, out\rangle$.

We note that one could also consider H_0^{ren} operators which depend on ϵ . Such a dependence can have an effect on the phase and normalization of $\lim_{\epsilon \rightarrow +0} \lim_{T \rightarrow \infty} \hat{U}_\epsilon^{\text{ren}}(0, -T) | v_i \rangle$ and $\lim_{\epsilon \rightarrow +0} \lim_{T \rightarrow \infty} \hat{U}_\epsilon^{\text{ren}}(0, T) | v_i \rangle$. In particular, one can take ϵ -dependent renormalized energies in (15). The requirement (18) is relaxed then to

$$\lim_{\epsilon \rightarrow +0} E_i^{\text{ren}}(\epsilon) = E_i. \quad (24)$$

It is not difficult to check that taking a renormalized energy level that depends on ϵ as

$$E_i^{\text{ren}}(\epsilon) = E_i + i\hbar\epsilon c_i + O(\epsilon^2), \quad (25)$$

where c_i is a complex number, will change the normalization of $\lim_{\epsilon \rightarrow +0} \lim_{T \rightarrow \infty} \hat{U}_\epsilon^{\text{ren}}(0, -T) | v_i \rangle$ by a factor

$$\frac{1}{1 + c_i} \quad (26)$$

and the normalization of $\lim_{\epsilon \rightarrow +0} \lim_{T \rightarrow \infty} \hat{U}_\epsilon^{\text{ren}}(0, T) | v_i \rangle$ by a factor

$$\frac{1}{1 - c_i}. \quad (27)$$

If c_i is real, then these factors are also real, but for a complex c_i they have a nonzero phase as well. In this paper we restrict ourselves to H_0^{ren} operators which are independent of ϵ .

3.3 Regularization and further remarks

The expressions (22) and (23) are somewhat formal, since $|v^{\text{ren}}\rangle$ is usually not a normalizable eigenvector, thus the denominator is not a finite number. If necessary, one can introduce a regularization to properly define these quotients. One possible regularization method is the following: one takes superpositions

$$|v_i^{\text{ren}}, \mu\rangle = \int da f_\mu(i, a) |v_a^{\text{ren}}\rangle, \quad (28)$$

where μ is a regularization parameter and $f_\mu(i, a)$ is a suitable function with the property that $\langle v_i^{\text{ren}}, \mu | v_j^{\text{ren}}, \mu \rangle$ is finite for all i and j if $\mu \neq 0$. The original $|v_i^{\text{ren}}\rangle$ vectors should be recovered in the $\mu \rightarrow 0$ limit. In the formulas (20), (21), (22) and (23) the vectors $|v^{\text{ren}}\rangle$ should be replaced by $|v^{\text{ren}}, \mu\rangle$.

Quantum field theories usually exhibit a type of divergence that is related to the infiniteness of the volume of the space. In particular, the powers of H_{int} are not well defined, because these powers involve vacuum bubble diagrams, and the product of the Dirac-deltas expressing momentum conservation in the vertices of a vacuum bubble diagram is not well defined. In order to deal with this divergence one has to introduce a suitable regularization of the powers of the H_{int} operator, and thus of the time evolution operator. The parameter for this regularization will be denoted by λ . H_0^{ren} has to be taken to depend on λ , and it will not have a finite limit as $\lambda \rightarrow 0$, because the eigenvalues of H are divergent. All eigenvalues of H contain the same additive divergent part, therefore this type of divergence cancels out in the differences of the eigenvalues. An illustration will be given in section 5.

Taking into consideration the regularizations described above, the formulas (20), (21), (22) and (23) take the following form:

$$|v, in\rangle = \lim_{\lambda \rightarrow 0} \lim_{\epsilon \rightarrow +0} \lim_{\mu \rightarrow 0} \lim_{T \rightarrow \infty} \frac{1}{\sqrt{Z_{v, \lambda, \epsilon, \mu, in}(-T)}} \hat{U}_\epsilon^{\text{ren}}(0, -T) |v^{\text{ren}}, \mu\rangle_\lambda, \quad (29)$$

$$|v, out\rangle = \lim_{\lambda \rightarrow 0} \lim_{\epsilon \rightarrow +0} \lim_{\mu \rightarrow 0} \lim_{T \rightarrow \infty} \frac{1}{\sqrt{Z_{v, \lambda, \epsilon, \mu, out}(T)}} \hat{U}_\epsilon^{\text{ren}}(0, T) |v^{\text{ren}}, \mu\rangle_\lambda, \quad (30)$$

$$Z_{v, \lambda, \epsilon, \mu, in}(-T) = \frac{\langle v^{\text{ren}}, \mu | \hat{U}_\epsilon^{\text{ren}}(0, -T)^\dagger \hat{U}_\epsilon^{\text{ren}}(0, -T) | v^{\text{ren}}, \mu \rangle_\lambda}{\langle v^{\text{ren}}, \mu | v^{\text{ren}}, \mu \rangle}, \quad (31)$$

$$Z_{v, \lambda, \epsilon, \mu, out}(T) = \frac{\langle v^{\text{ren}}, \mu | \hat{U}_\epsilon^{\text{ren}}(0, T)^\dagger \hat{U}_\epsilon^{\text{ren}}(0, T) | v^{\text{ren}}, \mu \rangle_\lambda}{\langle v^{\text{ren}}, \mu | v^{\text{ren}}, \mu \rangle}. \quad (32)$$

If one calculates a quantity involving *in* or *out* states, then generally the calculation should be done with nonzero λ , ϵ and μ , and the $\mu \rightarrow 0$, $\epsilon \rightarrow +0$ and $\lambda \rightarrow 0$ limits should be taken at the end. For instance, one should calculate the matrix element $\langle v_i, in | O | v_j, in \rangle$, where O is an operator, as $\lim_{\lambda \rightarrow 0} \lim_{\epsilon \rightarrow +0} \lim_{\mu \rightarrow 0} \langle v_i, in, \epsilon, \mu | O | v_j, in, \epsilon, \mu \rangle_\lambda$. The regularization associated with λ should be done on the whole product $\langle v_i, in, \epsilon, \mu | O | v_j, in, \epsilon, \mu \rangle$, not separately on $|v_j, in, \epsilon, \mu\rangle$ and $|v_i, in, \epsilon, \mu\rangle$. This rule applies also when the S-matrix elements are calculated.

It is worth noting that in quantum mechanical potential scattering (assuming short range and sufficiently regular potentials) the self-energy corrections are usually 0 (thus the renormalization of H_0 is not necessary) and the normalization factors $Z_{v, in}$ and $Z_{v, out}$ are usually equal to 1 after the $T \rightarrow \infty$, $\mu \rightarrow 0$, $\epsilon \rightarrow +0$ and $\lambda \rightarrow 0$ limits are taken (in fact, it is not necessary to introduce the regularization to which the λ parameter

belongs). We refer the reader to [14], section 3, example 2 for a hint about the type of argument that can be used to derive these results. In quantum field theory, however, the energy renormalizations and the normalization constants usually have nontrivial values. This can be seen in specific examples (e.g. in the ϕ^4 model, see section 5) by perturbative calculations.

For a nondegenerate state $|v^{\text{ren}}\rangle$ the phase of $|v, in\rangle$ defined by (20) (or (29)) is such that $\langle v^{\text{ren}}|v, in\rangle$ is real, and the same can be said of $|v, out\rangle$. This implies that $|v, in\rangle = |v, out\rangle$ and thus $\langle v, out|v, in\rangle = \langle v^{\text{ren}}|v^{\text{ren}}\rangle$. In quantum field theory, in particular, the definitions (29) and (30) yield the result $S_{\Omega;\Omega} = 1$ for the S-matrix element describing the vacuum-vacuum transition. The one-particle S-matrix element for a stable particle will be $S_{k_1;k_2} = \langle k_1|k_2\rangle = \delta^3(k_1 - k_2)$, where k denotes the momentum of the particle.

3.4 Perturbation theory

In this section we discuss some aspects of the application of perturbation theory to (20) and (21). The perturbation series for $\lim_{T \rightarrow \infty} \langle v_i|\hat{U}_\epsilon(0, -T)|v_j\rangle$ and $\lim_{T \rightarrow \infty} \langle v_i|\hat{U}_\epsilon(0, T)^\dagger|v_j\rangle$ with unrenormalized H_0 serve as basic ingredients for this.

The perturbation series for $\lim_{T \rightarrow \infty} \langle v_i|\hat{U}_\epsilon(0, -T)|v_j\rangle$ with unrenormalized H_0 reads as

$$\begin{aligned} \lim_{T \rightarrow \infty} \langle v_i|\hat{U}_\epsilon(0, -T)|v_j\rangle &= \langle v_i|v_j\rangle + \frac{-i}{\hbar} g \frac{\langle ij\rangle}{P(ij) + \epsilon} \\ &+ \sum_{k=2}^{\infty} \left(\frac{-i}{\hbar} \right)^k g^k \int dm_1 dm_2 \dots dm_{k-1} \frac{\langle im_{k-1}\rangle}{P(ij) + \epsilon} \frac{\langle m_{k-1}m_{k-2}\rangle}{P(m_{k-1}j) + \epsilon} \dots \frac{\langle m_2m_1\rangle}{P(m_2j) + \epsilon} \frac{\langle m_1j\rangle}{P(m_1j) + \epsilon}, \end{aligned} \quad (33)$$

where the notation

$$P(ij) = \frac{i}{\hbar} (E_i^0 - E_j^0), \quad \langle ij\rangle = \langle v_i|\frac{H - H_0}{g}|v_j\rangle = \langle v_i|H_{\text{int}}|v_j\rangle \quad (34)$$

is used, and the integrations over the intermediate states are understood in a general sense, in particular they include summation over discrete states. E_i^0 denotes the eigenvalues of H_0 . The object $\langle ij\rangle$ is a generalized function of the variables i and j .

For $\lim_{T \rightarrow \infty} \langle v_i|\hat{U}_\epsilon(0, T)^\dagger|v_j\rangle$ we have

$$\begin{aligned} \lim_{T \rightarrow \infty} \langle v_i|\hat{U}_\epsilon(0, T)^\dagger|v_j\rangle &= \langle v_i|v_j\rangle + \frac{-i}{\hbar} g \frac{\langle ij\rangle}{P(ji) + \epsilon} \\ &+ \sum_{k=2}^{\infty} \left(\frac{-i}{\hbar} \right)^k g^k \int dm_1 dm_2 \dots dm_{k-1} \frac{\langle im_1\rangle}{P(m_1i) + \epsilon} \frac{\langle m_1m_2\rangle}{P(m_2i) + \epsilon} \dots \frac{\langle m_{k-2}m_{k-1}\rangle}{P(m_{k-1}i) + \epsilon} \frac{\langle m_{k-1}j\rangle}{P(ji) + \epsilon}. \end{aligned} \quad (35)$$

These formulas can be used as a basis for deriving perturbative formulas for the components of the *in* and *out* states, for the S-matrix elements and for other quantities.

In order to take into account the renormalization of H_0 one should replace in the formulas (33) and (35) the $P(ij)$ given in (34) by

$$P^{\text{ren}}(ij) = \frac{i}{\hbar}(E_i^{\text{ren}} - E_j^{\text{ren}}), \quad (36)$$

and the $\langle ij \rangle$ given in (34) by

$$\langle ij \rangle^{\text{ren}} = \langle v_i^{\text{ren}} | \frac{H - H_0^{\text{ren}}}{g} | v_j^{\text{ren}} \rangle. \quad (37)$$

In this way one obtains series for $\lim_{T \rightarrow \infty} \langle v_i^{\text{ren}} | \hat{U}_\epsilon^{\text{ren}}(0, -T) | v_j^{\text{ren}} \rangle$ and $\lim_{T \rightarrow \infty} \langle v_i^{\text{ren}} | \hat{U}_\epsilon^{\text{ren}}(0, T)^\dagger | v_j^{\text{ren}} \rangle$. It is assumed, of course, that the $|v_i^{\text{ren}}\rangle$ vectors constitute a basis and their normalization is $\langle v_i^{\text{ren}} | v_j^{\text{ren}} \rangle = \delta(i, j)$. Since H_0^{ren} depends on g , the series for $\lim_{T \rightarrow \infty} \langle v_i^{\text{ren}} | \hat{U}_\epsilon^{\text{ren}}(0, -T) | v_j^{\text{ren}} \rangle$ and $\lim_{T \rightarrow \infty} \langle v_i^{\text{ren}} | \hat{U}_\epsilon^{\text{ren}}(0, T)^\dagger | v_j^{\text{ren}} \rangle$ will not be power series in g ; in order to obtain power series one has to expand their terms and collect the terms having the same power of g .

We note that in the case when the Hilbert space is finite dimensional the Rayleigh-Schrödinger perturbation theory can be expected to be reproduced in effect. The integrations in (33) and (35) reduce to summations in this case. The divergences in the $\epsilon \rightarrow +0$ limit without the renormalization arise from those terms in which a zero energy difference $P(ab)$ occurs in a denominator.

We consider now the case when an eigenvector $|v_i\rangle$ of H_0 belongs to a degenerate eigenspace. Let us assume that H_0^{ren} takes the form (15). We find at first order that $|v_i\rangle$ must be such that $\langle ji \rangle = 0$ for all $j \neq i$ for which $E_j^0 = E_i^0$. This condition is the same as the condition that one finds in Rayleigh-Schrödinger perturbation theory at first order, and it is always possible to choose the basis vectors in the eigenspace of H_0 belonging to E_i^0 in such a way that this condition be satisfied. At second order we find that

$$\sum_m \frac{\langle jm \rangle \langle mi \rangle}{E_m^0 - E_i^0} = 0 \quad (38)$$

has to hold for all values of $j \neq i$ for which $E_j^0 = E_i^0$, where the summation has to be done over those values of m for which $E_m^0 \neq E_i^0$. This condition is stronger than the one in Rayleigh-Schrödinger perturbation theory; in the latter case (38) is required to hold only for those values of j for which $|v_j, in\rangle$ remains degenerate with $|v_i, in\rangle$ at first order. While the weaker condition of the Rayleigh-Schrödinger perturbation theory can always be satisfied by choosing suitable basis vectors, the stronger condition (38) cannot always be satisfied, without violating the condition at first order, if the degeneracy of the eigenspace belonging to E_i^0 is broken at first order. This shows that the form (15) for H_0^{ren} is not suitable for all cases.

Finally, it follows from the formulas (33) and (35) that the vectors $\lim_{T \rightarrow \infty} \hat{U}_\epsilon(0, -T)|v_j\rangle$ and $\lim_{T \rightarrow \infty} \hat{U}_\epsilon(0, T)|v_j\rangle$ satisfy the Lippmann-Schwinger equations

$$\lim_{T \rightarrow \infty} \hat{U}_\epsilon(0, -T)|v_j\rangle = |v_j\rangle + \frac{-i}{\hbar} g \frac{1}{\frac{i}{\hbar}(H_0 - E_j^0) + \epsilon} H_{\text{int}} \lim_{T \rightarrow \infty} \hat{U}_\epsilon(0, -T)|v_j\rangle, \quad (39)$$

$$\lim_{T \rightarrow \infty} \hat{U}_\epsilon(0, T)|v_j\rangle = |v_j\rangle + \frac{-i}{\hbar} g \frac{1}{\frac{i}{\hbar}(H_0 - E_j^0) - \epsilon} H_{\text{int}} \lim_{T \rightarrow \infty} \hat{U}_\epsilon(0, T)|v_j\rangle, \quad (40)$$

where $\epsilon > 0$. Similar equations can be written with H_0^{ren} ; these equations take the form

$$\lim_{T \rightarrow \infty} \hat{U}_\epsilon^{\text{ren}}(0, -T)|v_j\rangle = |v_j^{\text{ren}}\rangle + \frac{-i}{\hbar} \frac{1}{\frac{i}{\hbar}(H_0^{\text{ren}} - E_j^{\text{ren}}) + \epsilon} (H - H_0^{\text{ren}}) \lim_{T \rightarrow \infty} \hat{U}_\epsilon^{\text{ren}}(0, -T)|v_j^{\text{ren}}\rangle, \quad (41)$$

$$\lim_{T \rightarrow \infty} \hat{U}_\epsilon^{\text{ren}}(0, T)|v_j\rangle = |v_j^{\text{ren}}\rangle + \frac{-i}{\hbar} \frac{1}{\frac{i}{\hbar}(H_0^{\text{ren}} - E_j^{\text{ren}}) - \epsilon} (H - H_0^{\text{ren}}) \lim_{T \rightarrow \infty} \hat{U}_\epsilon^{\text{ren}}(0, T)|v_j^{\text{ren}}\rangle. \quad (42)$$

4 The method of adiabatic switching

Another method that can be used to define the $T \rightarrow \infty$ limit of $U(0, -T)|v\rangle$ and $U(0, T)|v\rangle$, instead of the averaging procedure described in section 3, is the adiabatic switching of the interaction. In this case the *in* and *out* states are given by

$$|v, in\rangle = \lim_{\epsilon \rightarrow +0} \lim_{T \rightarrow \infty} U_\epsilon(0, -T)|v\rangle, \quad (43)$$

$$|v, out\rangle = \lim_{\epsilon \rightarrow +0} \lim_{T \rightarrow \infty} U_\epsilon(0, T)|v\rangle, \quad (44)$$

where

$$U_\epsilon(t_2, t_1) = T \exp \left[-\frac{i}{\hbar} \int_{t_1}^{t_2} g H_{\text{int}, \epsilon}(t) dt \right] \quad (45)$$

and

$$H_{\text{int}, \epsilon}(t) = e^{-\epsilon|t|} e^{\frac{i}{\hbar} H_0 t} H_{\text{int}} e^{-\frac{i}{\hbar} H_0 t}. \quad (46)$$

In (45) T denotes time ordering. In potential scattering this definition, like (7) and (8), is usually satisfactory. In general, however, $\lim_{T \rightarrow \infty} U_\epsilon(0, -T)$ and $\lim_{T \rightarrow \infty} U_\epsilon(0, T)$ are not convergent as $\epsilon \rightarrow +0$, and, correspondingly, the vectors $|v, in\rangle$ and $|v, out\rangle$ cannot be defined. Similarly to the case of the method of averaging, this divergence can be related to the existence of nonzero self-energy corrections.

4.1 Renormalization of H_0

The problem of divergence in the $\epsilon \rightarrow +0$ limit can be handled by the renormalization of H_0 , i.e. by a suitable choice for H_0^{ren} , which is used to define $U_\epsilon^{\text{ren}}(t_2, t_1)$ by substituting

$$e^{-\epsilon|t|} e^{\frac{i}{\hbar} H_0^{\text{ren}} t} (H - H_0^{\text{ren}}) e^{-\frac{i}{\hbar} H_0^{\text{ren}} t} \quad (47)$$

for $gH_{\text{int},\epsilon}(t)$ in (45). The definitions for $|v, in\rangle$ and $|v, out\rangle$ are thus

$$|v, in\rangle = \lim_{\epsilon \rightarrow +0} \lim_{T \rightarrow \infty} U_\epsilon^{\text{ren}}(0, -T) |v^{\text{ren}}\rangle, \quad (48)$$

$$|v, out\rangle = \lim_{\epsilon \rightarrow +0} \lim_{T \rightarrow \infty} U_\epsilon^{\text{ren}}(0, T) |v^{\text{ren}}\rangle. \quad (49)$$

The vector $|v^{\text{ren}}\rangle$ appearing in (48) and (49) should be an eigenvector of H_0^{ren} . The choice of H_0^{ren} is restricted by the requirement that the $\epsilon \rightarrow +0$ limits in these formulas should exist.

In general, the H_0^{ren} operators that are suitable for the method of adiabatic switching are different from those that are suitable for the method of averaging. In particular, it is not true that H_0^{ren} has to have the same eigenvalues as H . In order to illustrate this point we consider the case when the Hilbert space is finite dimensional and H_0 has nondegenerate spectrum. The H_0^{ren} operator is taken to be diagonal, i.e. of the form (15). It is not difficult to check, using the perturbation formulas (58) and (59) below, that if the adiabatic switching is applied, then the necessary shift of the eigenvalues of H_0 is not the full self-energy correction (i.e. the difference between the energy of $|v, in\rangle$ and the energy of $|v\rangle$). Using the notation $\delta E_j = g\delta E_j^1 + g^2\delta E_j^2 + g^3\delta E_j^3 + \dots$ for the self-energy correction (for a state $|v_j\rangle$) and $\Delta E_j = g\Delta E_j^1 + g^2\Delta E_j^2 + g^3\Delta E_j^3 + \dots$ for the required energy renormalization, we find that

$$\Delta E_j^1 = \delta E_j^1 \quad \text{and} \quad \Delta E_j^2 = \frac{1}{2}\delta E_j^2, \quad (50)$$

i.e. at first order the two quantities agree, but at second order the required energy renormalization is only the half of the self-energy correction. For ΔE_j^3 we obtain

$$\Delta E_j^3 = \sum_{m_1, m_2 \neq j; m_1 \neq m_2} \frac{\langle jm_2 \rangle \langle m_2 m_1 \rangle \langle m_1 j \rangle}{3(E_{m_2}^0 - E_j^0)(E_{m_1}^0 - E_j^0)} + \sum_{m \neq j} \frac{\langle jm \rangle \langle mj \rangle (\langle mm \rangle - \langle jj \rangle)}{2(E_m^0 - E_j^0)^2}, \quad (51)$$

where the notation of section 4.4 is used. The expression for δE_j^3 , that is well known from Rayleigh-Schrödinger perturbation theory, is

$$\delta E_j^3 = \sum_{m_1, m_2 \neq j; m_1 \neq m_2} \frac{\langle jm_2 \rangle \langle m_2 m_1 \rangle \langle m_1 j \rangle}{(E_{m_2}^0 - E_j^0)(E_{m_1}^0 - E_j^0)} + \sum_{m \neq j} \frac{\langle jm \rangle \langle mj \rangle (\langle mm \rangle - \langle jj \rangle)}{(E_m^0 - E_j^0)^2}. \quad (52)$$

This shows that there is no simple relation between ΔE_j^3 and δE_j^3 .

Although the energy renormalizations are generally not equal to the self-energy corrections, the latter can of course be obtained from the eigenvalue equations for the *in* and *out* states.

4.2 Normalization of the *in* and *out* states, regularization

Another significant difference from the method of averaging is that $\lim_{T \rightarrow \infty} U_\epsilon^{\text{ren}}(0, -T)$, $\lim_{T \rightarrow \infty} U_\epsilon^{\text{ren}}(0, T)$ and their limits as $\epsilon \rightarrow +0$ are unitary (more precisely, scalar product preserving), therefore it is not necessary to include any normalization factor in (43) and (44).

If one takes renormalized energy levels in (15) that depend on ϵ , then the phase of the *in* and *out* states can change. In this paper we restrict ourselves to H_0^{ren} operators that do not depend on ϵ .

Regularizations similar to those described in section 3.3 might be needed also in the case of the method of adiabatic switching. Taking into account these regularizations, the formulas (43) and (44) have the form

$$|v, in\rangle = \lim_{\lambda \rightarrow 0} \lim_{\epsilon \rightarrow +0} \lim_{\mu \rightarrow 0} \lim_{T \rightarrow \infty} U_\epsilon^{\text{ren}}(0, -T) |v^{\text{ren}}, \mu\rangle_\lambda, \quad (53)$$

$$|v, out\rangle = \lim_{\lambda \rightarrow 0} \lim_{\epsilon \rightarrow +0} \lim_{\mu \rightarrow 0} \lim_{T \rightarrow \infty} U_\epsilon^{\text{ren}}(0, T) |v^{\text{ren}}, \mu\rangle_\lambda. \quad (54)$$

4.3 Modification by phase factors

In quantum field theory the use of the method of adiabatic switching seems to be restricted mainly to the vacuum state, and in this case the Gell-Mann–Low formula [22, 23] is used, which does not require the renormalization of H_0 .

In [14] we proposed the following formulas, which can be regarded as generalizations of the Gell-Mann–Low formula:

$$|v, in\rangle = \lim_{\epsilon \rightarrow +0} \lim_{T \rightarrow \infty} \Pi_{v,\epsilon}(-T) U_\epsilon(0, -T) |v\rangle, \quad (55)$$

$$|v, out\rangle = \lim_{\epsilon \rightarrow +0} \lim_{T \rightarrow \infty} \Pi_{v,\epsilon}(T) U_\epsilon(0, T) |v\rangle, \quad (56)$$

where $\Pi_{v,\epsilon}(T)$ is a complex number of absolute value 1, i.e. a phase factor, given by

$$\Pi_{v,\epsilon}(T) = \frac{\sqrt{\langle v | U_\epsilon(T, 0) | v \rangle \langle v | U_\epsilon(0, T) | v \rangle}}{\langle v | U_\epsilon(0, T) | v \rangle}. \quad (57)$$

It may, of course, be necessary to introduce regularizations similar to those described in section 3.3. The phase factors in (55) and (56) have no effect on transition probabilities. In [14] we stated the conjecture that in a large class of quantum field theories the divergence in the $\epsilon \rightarrow +0$ limit is entirely canceled out by the phase factors, without any need for the renormalization of H_0 . Nevertheless, the renormalization of H_0 is needed in some cases also with (55) and (56).

4.4 Perturbation theory

The perturbation series for $\lim_{T \rightarrow \infty} \langle v_i | U_\epsilon(0, -T) | v_j \rangle$ with unrenormalized H_0 reads as

$$\begin{aligned} \lim_{T \rightarrow \infty} \langle v_i | U_\epsilon(0, -T) | v_j \rangle = & \langle v_i | v_j \rangle + \frac{-i}{\hbar} g \frac{\langle ij \rangle}{P(ij) + \epsilon} + \sum_{k=2}^{\infty} \left(\frac{-i}{\hbar} \right)^k g^k \int dm_1 dm_2 \dots dm_{k-1} \\ & \frac{\langle im_{k-1} \rangle}{P(ij) + k\epsilon} \frac{\langle m_{k-1} m_{k-2} \rangle}{P(m_{k-1}j) + (k-1)\epsilon} \dots \frac{\langle m_2 m_1 \rangle}{P(m_2j) + 2\epsilon} \frac{\langle m_1 j \rangle}{P(m_1j) + \epsilon}, \end{aligned} \quad (58)$$

where the notation (34) is used. This formula differs from (33) in the coefficients of ϵ in the denominators. To first order, however, there is no difference between the method of adiabatic switching and the method of averaging. This implies, in particular, that the Z normalization constants in (20) and (21) can be taken to be 1 to first order.

For $\lim_{T \rightarrow \infty} \langle v_i | U_\epsilon(T, 0) | v_j \rangle$ we have

$$\begin{aligned} \lim_{T \rightarrow \infty} \langle v_i | U_\epsilon(T, 0) | v_j \rangle = & \langle v_i | v_j \rangle + \frac{-i}{\hbar} g \frac{\langle ij \rangle}{P(ji) + \epsilon} + \sum_{k=2}^{\infty} \left(\frac{-i}{\hbar} \right)^k g^k \int dm_1 dm_2 \dots dm_{k-1} \\ & \frac{\langle im_1 \rangle}{P(m_1i) + \epsilon} \frac{\langle m_1 m_2 \rangle}{P(m_2i) + 2\epsilon} \dots \frac{\langle m_{k-2} m_{k-1} \rangle}{P(m_{k-1}i) + (k-1)\epsilon} \frac{\langle m_{k-1} j \rangle}{P(ji) + k\epsilon}. \end{aligned} \quad (59)$$

These formulas can be used as a basis for deriving perturbative formulas for the components of the *in* and *out* states, for the S-matrix elements and for other quantities. The renormalization of H_0 should be taken into consideration as described in section 3.4.

Concerning the suitability of (15) and the choice of the basis vectors in degenerate eigenspaces of H_0 in the case of finite dimensional Hilbert spaces, one finds similar results as in the case of averaging.

A remarkable consequence of the fact that the coefficients of ϵ in the denominators of (58), (59) and in (33), (35), respectively, are different, is that the vectors $\lim_{T \rightarrow \infty} U_\epsilon(0, -T) | v_j \rangle$ and $\lim_{T \rightarrow \infty} U_\epsilon(0, T) | v_j \rangle$ do not satisfy Lippmann-Schwinger equations similar to (39) and (40). Similar statement can be made about $\lim_{T \rightarrow \infty} U_\epsilon^{\text{ren}}(0, -T) | v_j^{\text{ren}} \rangle$ and $\lim_{T \rightarrow \infty} U_\epsilon^{\text{ren}}(0, T) | v_j^{\text{ren}} \rangle$.

5 Illustration: ϕ^4 model

In this section we illustrate the ideas described in sections 3 and 4 in the case of the ϕ^4 model. We discuss the vacuum-vacuum S-matrix element $S_{\Omega, \Omega}$, the one-particle S-matrix element $S_{k_1; k_2}$ and the two-particle S-matrix element $S_{k_3 k_4; k_1 k_2}$. We calculate these S-matrix elements using perturbation theory to second order, and some third order terms will also be considered in the case of $S_{k_3 k_4; k_1 k_2}$. The results are ‘bare’, i.e. they are expressed in terms of the initial unrenormalized parameters of the full Hamiltonian operator

H . The ϕ^4 model is known to contain ultraviolet divergences. These are not relevant for the subject of this paper, nevertheless they can be avoided by introducing a momentum cutoff. The calculations are straightforward, but rather lengthy, therefore we shall not write many details of them. Basic formulas concerning the ϕ^4 model, which are relevant for the calculations, are presented in Appendix A. We take H_0^{ren} to be of the form (15).

5.1 The method of averaging

5.1.1 The vacuum-vacuum S-matrix element

Without energy renormalization we obtain to second order in g that

$$\begin{aligned} & \lim_{T \rightarrow \infty} \langle \Omega | \hat{U}_\epsilon(0, T)^\dagger \hat{U}_\epsilon(0, -T) | \Omega \rangle_\lambda = \\ & = 1 + \left(\frac{-i}{\hbar} \right)^2 g^2 \int \frac{d^3 k_A d^3 k_B d^3 k_C d^3 k_D}{4!} \frac{\eta^2}{\omega_A \omega_B \omega_C \omega_D} \frac{2[\delta_\lambda^3(k_A + k_B + k_C + k_D)]^2}{\epsilon \left[\frac{i}{\hbar}(\omega_A + \omega_B + \omega_C + \omega_D) + \epsilon \right]} \\ & \quad + \left(\frac{-i}{\hbar} \right)^2 g^2 \int \frac{d^3 k_A d^3 k_B d^3 k_C d^3 k_D}{4!} \frac{\eta^2}{\omega_A \omega_B \omega_C \omega_D} \frac{[\delta_\lambda^3(k_A + k_B + k_C + k_D)]^2}{\left[\frac{i}{\hbar}(\omega_A + \omega_B + \omega_C + \omega_D) + \epsilon \right]^2}, \end{aligned} \quad (60)$$

where $\eta = (2\pi)^3 \frac{4!}{2^{12}\pi^6}$. This quantity diverges as $1/\epsilon$ in the $\epsilon \rightarrow +0$ limit and as $1/\lambda^3$ in the $\lambda \rightarrow 0$ limit. (The λ parameter is explained at the end of Appendix A.)

As explained in section 3, energy renormalization and normalization factors are needed to turn (60) into the correct vacuum-vacuum S-matrix element. The energy renormalization for $|\Omega\rangle$ is denoted as $\Delta E_\Omega = g\Delta E_\Omega^1 + g^2\Delta E_\Omega^2 + \dots$. We restrict ourselves to a ΔE_Ω that does not depend on ϵ . It is straightforward to see that ΔE_Ω^1 has to be 0 in order to get a finite result for $\lim_{T \rightarrow \infty} \langle \Omega | \hat{U}_\epsilon^{\text{ren}}(0, T)^\dagger \hat{U}_\epsilon^{\text{ren}}(0, -T) | \Omega \rangle_\lambda$ in the $\epsilon \rightarrow +0$ limit. Calculating to second order we obtain then that

$$\lim_{T \rightarrow \infty} \langle \Omega | \hat{U}_\epsilon^{\text{ren}}(0, T)^\dagger \hat{U}_\epsilon^{\text{ren}}(0, -T) | \Omega \rangle_\lambda = \lim_{T \rightarrow \infty} \langle \Omega | \hat{U}_\epsilon(0, T)^\dagger \hat{U}_\epsilon(0, -T) | \Omega \rangle_\lambda + g^2 \frac{i}{\hbar} \frac{2\Delta E_\Omega^2}{\epsilon}. \quad (61)$$

From comparison with (60) one can see that ΔE_Ω^2 has to be chosen as

$$\Delta E_\Omega^2(\lambda) = - \int \frac{d^3 k_A d^3 k_B d^3 k_C d^3 k_D}{4!} \frac{\eta^2}{\omega_A \omega_B \omega_C \omega_D} \frac{[\delta_\lambda^3(k_A + k_B + k_C + k_D)]^2}{\omega_A + \omega_B + \omega_C + \omega_D} \quad (62)$$

in order for $\lim_{T \rightarrow \infty} \langle \Omega | \hat{U}_\epsilon^{\text{ren}}(0, T)^\dagger \hat{U}_\epsilon^{\text{ren}}(0, -T) | \Omega \rangle_\lambda$ to be finite when $\epsilon \rightarrow +0$. $\Delta E_\Omega^2(\lambda)$ diverges as $1/\lambda^3$ in the $\lambda \rightarrow 0$ limit.

For the constants $Z_{\Omega, \epsilon, \lambda, \text{in}}$ and $Z_{\Omega, \epsilon, \lambda, \text{out}}$ we obtain

$$\begin{aligned} Z_{\Omega, \epsilon, \lambda, \text{in}} &= 1 + \left(\frac{-i}{\hbar} \right)^2 g^2 \int \frac{d^3 k_A d^3 k_B d^3 k_C d^3 k_D}{4!} \frac{\eta^2}{\omega_A \omega_B \omega_C \omega_D} \\ &\quad \times \frac{[\delta_\lambda^3(k_A + k_B + k_C + k_D)]^2}{\left[\frac{i}{\hbar}(\omega_A + \omega_B + \omega_C + \omega_D) + \epsilon \right] \left[-\frac{i}{\hbar}(\omega_A + \omega_B + \omega_C + \omega_D) + \epsilon \right]} \end{aligned} \quad (63)$$

and $Z_{\Omega,\epsilon,\lambda,in} = Z_{\Omega,\epsilon,\lambda,out}$. These quantities are finite as $\epsilon \rightarrow +0$ but diverge as $1/\lambda^3$ in the $\lambda \rightarrow 0$ limit. We note that the energy renormalization has no effect on the $Z_{\Omega,\epsilon,\lambda,in}$ and $Z_{\Omega,\epsilon,\lambda,out}$ constants to second order in g .

Using the results (60)-(63) in the formulas (1) and (29)-(32) we find that the final result for the vacuum-vacuum S-matrix element to second order in g is $S_{\Omega,\Omega} = \langle \Omega, out | \Omega, in \rangle = 1$, in agreement with the remarks at the end of section 3.3.

5.1.2 The one-particle S-matrix element

To second order in g the contributions to $\lim_{T \rightarrow \infty} \langle k_2 | \hat{U}_\epsilon(0, T)^\dagger \hat{U}_\epsilon(0, -T) | k_1 \rangle_\lambda$ correspond to the graphs shown in figure 1. The rate of divergence of the second order part is $1/\epsilon$ in the $\epsilon \rightarrow +0$ limit.

In order to turn $\lim_{T \rightarrow \infty} \langle k_2 | \hat{U}_\epsilon(0, T)^\dagger \hat{U}_\epsilon(0, -T) | k_1 \rangle_\lambda$ into the correct one-particle S-matrix element energy renormalization and normalization factors are needed. From the requirement that the $\epsilon \rightarrow +0$ limit should be finite we obtain that the first order energy renormalization ΔE_k^1 has to be zero. The same condition yields that the second order energy renormalization has to be

$$\begin{aligned} \Delta E_k^2(\lambda) = & - \int \frac{d^3 k_A d^3 k_B}{3!} \frac{\eta^2}{\omega_A \omega_B \omega_C \omega (\omega_A + \omega_B + \omega_C - \omega)} \\ & - \int \frac{d^3 k_A d^3 k_B}{3!} \frac{\eta^2}{\omega_A \omega_B \omega_C \omega (\omega_A + \omega_B + \omega_C + \omega)} \\ & + \Delta E_\Omega^2(\lambda), \end{aligned} \quad (64)$$

where $k_C = k - k_A - k_B$ in ω_C and $\omega \equiv \omega(k)$.

$\Delta E_k^2(\lambda)$ diverges as $\lambda \rightarrow 0$, but the difference $\Delta E_k^2(\lambda) - \Delta E_\Omega^2(\lambda)$ has a finite limit (in fact, it is a constant function of λ).

For the constants $Z_{k,\epsilon,\lambda,in}$ and $Z_{k,\epsilon,\lambda,out}$ we obtain

$$\begin{aligned} Z_{k,\epsilon,\lambda,in} = Z_{k,\epsilon,\lambda,out} = & Z_{\Omega,\epsilon,\lambda,in} + \\ & + \left(\frac{-i}{\hbar} \right)^2 g^2 \int \frac{d^3 k_A d^3 k_B}{3!} \frac{\eta^2}{\omega_A \omega_B \omega_C \omega} \\ & \times \frac{1}{\left[\frac{i}{\hbar} (\omega_A + \omega_B + \omega_C - \omega) + \epsilon \right] \left[-\frac{i}{\hbar} (\omega_A + \omega_B + \omega_C - \omega) + \epsilon \right]} \\ & + \left(\frac{-i}{\hbar} \right)^2 g^2 \int \frac{d^3 k_A d^3 k_B}{3!} \frac{\eta^2}{\omega_A \omega_B \omega_C \omega} \\ & \times \frac{1}{\left[\frac{i}{\hbar} (\omega_A + \omega_B + \omega_C + \omega) + \epsilon \right] \left[-\frac{i}{\hbar} (\omega_A + \omega_B + \omega_C + \omega) + \epsilon \right]}. \end{aligned} \quad (65)$$

This quantity is finite in the $\epsilon \rightarrow +0$ limit and the energy renormalization does not have any effect on it. After these normalization factors and the energy renormalization

are taken into account we find that the result for the one-particle S-matrix element is $\langle k_2, out | k_1, in \rangle = \delta^3(k_2 - k_1)$, in accordance with the remarks at the end of section 3.3.

The energy renormalization $\lim_{\lambda \rightarrow 0} (\Delta E_k^2(\lambda) - \Delta E_\Omega^2(\lambda))$ is equal to the energy correction that can be obtained in the Lehmann-Symanzik-Zimmermann (LSZ) formalism. We note that both $Z_{k,\epsilon,\lambda,in}/Z_{\Omega,\epsilon,\lambda,in}$ and $\Delta E_k^2(\lambda) - \Delta E_\Omega^2(\lambda)$ are independent of λ .

5.1.3 The two-particle S-matrix element

In this section we consider the two-particle S-matrix element $\langle k_3, k_4, out | k_1, k_2, in \rangle$. The diagrams corresponding to the contributions to $\lim_{T \rightarrow \infty} \langle k_3, k_4 | \hat{U}_\epsilon(0, T)^\dagger \hat{U}_\epsilon(0, -T) | k_1, k_2 \rangle_\lambda$ to second order are shown in figure 2. There are additional graphs to be included in the calculation which are obtained by interchanging k_3 and k_4 in 2a, 2d, 2e, 2f and 2g; these are not shown in order to save space, and we will not mention them explicitly in the following. Those terms which are divergent in the $\epsilon \rightarrow +0$ limit correspond to the graphs 2e, 2f and 2g. The rate of divergence is $1/\epsilon$.

We find that the energy renormalization at second order is

$$\Delta E_{k_1 k_2}^2(\lambda) = \Delta E_\Omega^2(\lambda) + [\Delta E_{k_1}^2(\lambda) - \Delta E_\Omega^2(\lambda)] + [\Delta E_{k_2}^2(\lambda) - \Delta E_\Omega^2(\lambda)], \quad (66)$$

i.e. the energy renormalization for the two-particle state is the sum of the energy renormalizations for the constituent particles, if the vacuum energy renormalization is subtracted. This is in agreement with the natural expectation. It is also natural to expect that one would find a similar result to all orders of perturbation theory and for any number of particles.

For the normalization constants $Z_{k_1 k_2, \epsilon, \lambda, in}$ and $Z_{k_1 k_2, \epsilon, \lambda, out}$ we find that

$$Z_{k_1 k_2, \epsilon, \lambda, in} = Z_{k_1 k_2, \epsilon, \lambda, out} \quad (67)$$

and $Z_{k_1 k_2, \epsilon, \lambda, in}$ is given by the relation

$$\frac{Z_{k_1 k_2, \epsilon, \lambda, in}}{Z_{\Omega, \epsilon, \lambda, in}} = \frac{Z_{k_1, \epsilon, \lambda, in}}{Z_{\Omega, \epsilon, \lambda, in}} \frac{Z_{k_2, \epsilon, \lambda, in}}{Z_{\Omega, \epsilon, \lambda, in}}. \quad (68)$$

We conjecture that this relation holds to all orders of perturbation theory, and a similar relation holds for states containing arbitrary number of particles.

The joint effect of the energy renormalization and the normalization constants is that the contributions corresponding to the disconnected graphs 2e, 2f and 2g are entirely canceled out from the expression for the two-particle S-matrix, and only the terms corresponding to the graphs 2a, 2b, 2c and 2d remain.

At third order we restrict ourselves to the brief discussion of a certain type of graph, shown in figure 3, which describes interaction between the two particles and which contains a radiative correction on an external line. The terms corresponding to this graph are present in the expression for $\lim_{T \rightarrow \infty} \langle k_3, k_4 | \hat{U}_\epsilon(0, T)^\dagger \hat{U}_\epsilon(0, -T) | k_1, k_2 \rangle_\lambda$, and some of these




Figure 1: *Graphs for the one-particle S -matrix element to second order*




Figure 2: *Graphs for the two-particle S -matrix elements to second order*

terms are divergent as $1/\epsilon$ in the $\epsilon \rightarrow +0$ limit. The energy renormalization, in particular the modification of the matrix elements of H_{int} in (33) and (35) due to the second order part of the energy renormalization, eliminates this divergence.

At third order the modification of the denominators in (33) and (35) due to the energy renormalization is also relevant. This has the effect that the Dirac-delta factor $\delta(E_{k_3}^0 + E_{k_4}^0 - E_{k_1}^0 - E_{k_2}^0)$ in the contribution corresponding to the graph shown in figure 2b that expresses energy conservation becomes $\delta(E_{k_3} + E_{k_4} - E_{k_1} - E_{k_2})$, i.e. the unperturbed energy levels are replaced by the proper ones.




Figure 3: A third order graph containing a radiative correction on an external line

5.2 The method of adiabatic switching

In order to get the vacuum-vacuum S-matrix element we start with the matrix element $\lim_{T \rightarrow \infty} \langle \Omega | U_\epsilon(0, T)^\dagger U_\epsilon(0, -T) | \Omega \rangle_\lambda$, for which we obtain to second order in g that

$$\begin{aligned} & \lim_{T \rightarrow \infty} \langle \Omega | U_\epsilon(0, T)^\dagger U_\epsilon(0, -T) | \Omega \rangle_\lambda = \\ & = 1 + \left(\frac{-i}{\hbar} \right)^2 g^2 \int \frac{d^3 k_A d^3 k_B d^3 k_C d^3 k_D}{4!} \frac{\eta^2}{\omega_A \omega_B \omega_C \omega_D} \frac{[\delta_\lambda^3(k_A + k_B + k_C + k_D)]^2}{\epsilon [\frac{i}{\hbar}(\omega_A + \omega_B + \omega_C + \omega_D) + \epsilon]} \\ & + \left(\frac{-i}{\hbar} \right)^2 g^2 \int \frac{d^3 k_A d^3 k_B d^3 k_C d^3 k_D}{4!} \frac{\eta^2}{\omega_A \omega_B \omega_C \omega_D} \frac{[\delta_\lambda^3(k_A + k_B + k_C + k_D)]^2}{[\frac{i}{\hbar}(\omega_A + \omega_B + \omega_C + \omega_D) + \epsilon]^2}. \quad (69) \end{aligned}$$

This differs from (60), obtained in the case of the method of averaging, in a factor of 2 in the first integral on the right hand side.

Similarly to (61), we have

$$\lim_{T \rightarrow \infty} \langle \Omega | U_\epsilon^{\text{ren}}(0, T)^\dagger U_\epsilon^{\text{ren}}(0, -T) | \Omega \rangle_\lambda = \lim_{T \rightarrow \infty} \langle \Omega | U_\epsilon(0, T)^\dagger U_\epsilon(0, -T) | \Omega \rangle_\lambda + g^2 \frac{i}{\hbar} \frac{2\Delta E_\Omega^2}{\epsilon}. \quad (70)$$

The second term on the right hand side is of the same form as in (61). As a consequence, the second order energy renormalization needed in the case of adiabatic switching is 1/2 of that needed for the case of the averaging, in accordance with the discussion in section 4.1. Similar result can be found also for the one- and two-particle states. As we mentioned in section 4.2, normalization constants for the *in* and *out* states are not needed. The final results for the vacuum-vacuum, one-particle and two-particle S-matrix elements to second order are the same as in the case of the averaging.

Similarly to the case of averaging, the contribution of the graph 3 to $\lim_{T \rightarrow \infty} \langle \Omega | U_\epsilon(0, T)^\dagger U_\epsilon(0, -T) | \Omega \rangle_\lambda$ contains terms which are divergent as $1/\epsilon$ in the $\epsilon \rightarrow +0$ limit, but the energy renormalization, in particular the modification of the matrix elements of H_{int} due to the second order part of the energy renormalization, eliminates this divergence.

As regards the formulas (55) and (56) for the *in* and *out* states, we refer the reader to [14] for their discussion and their application to the ϕ^4 model. In the case of these

formulas those are the factors $\Pi_{v,\epsilon}$ (see (57)) that have the same effect as the energy renormalization has in the case of (48) and (49).

6 Conclusion

We discussed the methods of averaging and adiabatic switching for the definition of the infinite time limit of the time evolution operator $U(0, \pm T)$ in scattering theory. We described three significant differences between the two methods. The first difference is that the operator obtained using the method of averaging is not unitary in general, therefore normalization factors are needed in the definition of the *in* and *out* states in order to ensure the conservation of probabilities. In the case of the method of adiabatic switching such a problem of unitarity does not arise. The second difference is that in the case of the method of averaging the energy renormalizations needed in the free Hamiltonian operator are equal to the energy shifts caused by the interaction, whereas in the case of the adiabatic switching these quantities are generally different. The third difference between the two methods is that Lippmann-Schwinger equations exist in the case of the method of averaging, but similar equations do not hold in the case of the method of adiabatic switching. At the level of perturbation series these differences are encoded entirely in the (seemingly minor) difference between the coefficients of ϵ in the formulas (33), (35) and (58), (59). We presented an illustration of the application of the formulas for the *in* and *out* states in the case of the ϕ^4 model using perturbation theory to low orders. The final results that we obtained for the S-matrix elements were the same for the two methods, and they are in agreement with the expectations. It would be interesting to extend the illustration to all orders of perturbation theory and to other models. It would also be interesting to find general formulas for the energy renormalizations in the case of the method of adiabatic switching.

Appendix A

The ϕ^4 model has the Hamiltonian operator $H = H_0 + gH_{\text{int}}$, where H_0 is the Hamiltonian operator of the free scalar boson of mass m_0 :

$$H_0 = \frac{1}{2} \int d^3x [(\partial_t \phi)^2 + (\partial_x \phi)^2 + m_0^2 \phi^2] \quad (71)$$

and the interaction term H_{int} is

$$H_{\text{int}} = \int d^3x : \phi^4 : . \quad (72)$$

The free boson field is

$$\phi(x, t) = \int \frac{d^3k}{(2\sqrt{\pi})^3} [e^{ikx} e^{i\omega(k)t} a^\dagger(k) + e^{-ikx} e^{-i\omega(k)t} a(k)], \quad (73)$$

where

$$\omega(\mathbf{k}) = \sqrt{m_0^2 + \mathbf{k}^2}. \quad (74)$$

The particle annihilation operators $a(\mathbf{k})$ satisfy the commutation relation

$$[a(\mathbf{k}), a(\mathbf{k}')^\dagger] = \delta^3(\mathbf{k} - \mathbf{k}') \frac{1}{\omega(\mathbf{k})}. \quad (75)$$

For the boson field we have the commutation relation

$$[\phi(\mathbf{x}, t), \partial_t \phi(\mathbf{x}', t)] = i\delta^3(\mathbf{x} - \mathbf{x}'). \quad (76)$$

The multi-particle eigenstates of H_0 are

$$|k_1, k_2, \dots, k_n\rangle = \left(\prod_{i=1}^n \sqrt{\omega(k_i)} a(k_i)^\dagger \right) |\Omega\rangle, \quad (77)$$

where $|\Omega\rangle$ is the vacuum state of H_0 . These states have the normalization

$$\langle \Omega | \Omega \rangle = 1 \quad (78)$$

and

$$\langle k_1, k_2, \dots, k_n | k'_1, k'_2, \dots, k'_n \rangle = \sum_P \delta^3(k_1 - k_{P(1)}) \delta^3(k_2 - k_{P(2)}) \dots \delta^3(k_n - k_{P(n)}), \quad (79)$$

where the summation is done over the permutations of n elements.

The eigenvalues of H_0 for $|k_1, k_2, \dots, k_n\rangle$ and $|\Omega\rangle$ are

$$E_{k_1 k_2 \dots k_n}^0 = \omega(k_1) + \omega(k_2) + \dots + \omega(k_n) \quad (80)$$

and

$$E_\Omega^0 = 0. \quad (81)$$

It is straightforward to calculate the matrix elements of H_{int} ; a few examples read as

$$\langle k_1, k_2, k_3, k_4 | \int d^3\mathbf{x} : \phi^4 : |\Omega\rangle = (2\pi)^3 \delta^3(\mathbf{k}_1 + \mathbf{k}_2 + \mathbf{k}_3 + \mathbf{k}_4) \frac{1}{2^{12}\pi^6} \frac{4!}{\sqrt{\omega_1 \omega_2 \omega_3 \omega_4}} \quad (82)$$

$$\langle k_1, k_2, k_3 | \int d^3\mathbf{x} : \phi^4 : |k_4\rangle = (2\pi)^3 \delta^3(\mathbf{k}_1 + \mathbf{k}_2 + \mathbf{k}_3 - \mathbf{k}_4) \frac{1}{2^{12}\pi^6} \frac{4!}{\sqrt{\omega_1 \omega_2 \omega_3 \omega_4}} \quad (83)$$

$$\begin{aligned} \langle k_1, k_2, k_3, k_4, k_5 | \int d^3\mathbf{x} : \phi^4 : |k_6\rangle &= \langle k_1, k_2, k_3, k_4 | \int d^3\mathbf{x} : \phi^4 : |\Omega\rangle \delta^3(\mathbf{k}_5 - \mathbf{k}_6) \\ &+ \langle k_1, k_2, k_3, k_5 | \int d^3\mathbf{x} : \phi^4 : |\Omega\rangle \delta^3(\mathbf{k}_4 - \mathbf{k}_6) + \dots, \end{aligned} \quad (84)$$

where $\omega_i \equiv \omega(\mathbf{k}_i)$.

The regularization that is needed because of the divergence associated with the infiniteness of the volume of the space and that is mentioned in section 3.3 is introduced in the following way: if a diagram contains vacuum bubbles, then the Dirac-deltas expressing momentum conservation at the vertices in these vacuum bubbles are replaced by the regularized Dirac-deltas $\delta_\lambda^3(\mathbf{x}) = \delta_\lambda(x_1)\delta_\lambda(x_2)\delta_\lambda(x_3)$, where $\delta_\lambda(x) = 0$ if $x < -\lambda/2$ or $x > \lambda/2$ and $\delta_\lambda(x) = 1/\lambda$ if $-\lambda/2 \leq x \leq \lambda/2$.

Acknowledgments

This work was supported in part by the MIUR grant 2007JHLPEZ.

References

- [1] J. D. Kim, *Phys. Lett. B* **353** 213 (1995), arXiv:hep-th/9504018
- [2] J. D. Kim, arXiv:hep-th/9506031
- [3] N. Topor, *Mod. Phys. Lett. A* **12** 2951 (1997)
- [4] M. Perkins and P. Bowcock, *Nucl. Phys. B* **538** 612 (1999), arXiv:hep-th/9807146
- [5] E. Corrigan, *Int. J. Mod. Phys. A* **13** 2709 (1998), arXiv:hep-th/9707235
- [6] A. Chenaghlou, *Int. J. Mod. Phys. A* **15** 4623 (2000), arXiv:hep-th/0004121
- [7] A. Chenaghlou, *Int. J. Mod. Phys. A* **16** 4613 (2001), arXiv:hep-th/0108077
- [8] A. Chenaghlou and E. Corrigan, *Int. J. Mod. Phys. A* **15** 4417 (2000), arXiv:hep-th/0002065
- [9] M. Ablikim and E. Corrigan, *Int. J. Mod. Phys. A* **16** 625 (2001), arXiv:hep-th/0007214
- [10] M. Gattobigio, A. Liguori and M. Mintchev, *J. Math. Phys.* **40** 2949 (1999), arXiv:hep-th/9811188
- [11] G. Delfino, G. Mussardo and P. Simonetti, *Nucl. Phys. B* **473** 469 (1996)
- [12] Z. Bajnok, G. Bohm and G. Takacs, *J. Phys. A* **35** 9333 (2002), hep-th/0207079
- [13] L. Borda, L. Fritz, N. Andrei and G. Zarand, *Phys. Rev. B* **75** 235112 (2007), arXiv:cond-mat/0703293

- [14] G. Zs. Toth, *Cent. Eur. J. Phys.* (2009) DOI 10.2478/s11534-009-0139-0, arXiv:0812.3614 [hep-th]
- [15] J. R. Taylor, *Scattering theory: the quantum theory on nonrelativistic collisions*, Wiley, New York, NY, 1972
- [16] R. Newton, *Scattering theory of waves and particles*, 2nd ed., Springer-Verlag, Berlin, 1982
- [17] S. Weinberg, *The quantum theory of fields I*, Cambridge University Press, Cambridge, New York, 1995-2000
- [18] M. Gell-Mann and M. L. Goldberger, *Phys. Rev.* **91** 398 (1953)
- [19] M. E. Peskin, D. V. Schroeder, *An introduction to quantum field theory*, Addison-Wesley, Reading, Mass., 1995
- [20] J. Pirenne, *Helv. Phys. Acta* **21** 226 (1948)
- [21] J. Pirenne, *Phys. Rev.* **86** 395 (1952)
- [22] M. Gell-Mann and F. Low, *Phys. Rev.* **84** 350 (1951)
- [23] A. L. Fetter and J. D. Walecka, *Quantum theory of many-particle systems*, Dover Publications, Mineola, NY, 2003, chapter 3